

Appendix 8.1b - Phase 1 Habitats and principal NVC communities after additional 2014 survey

Phase 1 Habitat code	Phase 1 Habitat	Associated NVC communities	Hectares/% area
B1.1	Unimproved acid grassland	U4a <i>Festuca ovina</i> - <i>Agrostis capillaris</i> - <i>Galium saxatile</i> grassland, typical sub-community	5.74/1.22%
		U6 <i>Juncus squarrosus</i> - <i>Festuca ovina</i> grassland	
B5	Marsh/Marshy grassland	M25 <i>Molinia caerulea</i> - <i>Potentilla erecta</i> mire	0.71/0.15%
C1.1	Bracken	U20 <i>Pteridium aquilinum</i> - <i>Galium saxatile</i> community	0.84/0.18%
D1.1	Dry acid heath	H12a <i>Calluna vulgaris</i> - <i>Vaccinium myrtillus</i> heath, <i>Calluna vulgaris</i> sub-community	38.25/8.15%
		H10a <i>Calluna vulgaris</i> - <i>Erica cinerea</i> heath, typical sub-community	
D2	Wet heath	M15b <i>Scirpus cespitosus</i> - <i>Erica tetralix</i> wet heath, typical sub-community	149.74/31.93%
D5	Dry heath/acid grassland	H10a <i>Calluna vulgaris</i> - <i>Erica cinerea</i> heath, typical sub-community	1.43/0.30%
		H12a <i>Calluna vulgaris</i> - <i>Vaccinium myrtillus</i> heath, <i>Calluna vulgaris</i> sub-community	
		U4a <i>Nardus stricta</i> - <i>Galium saxatile</i> grassland, species-poor sub-community	
E1.6.1	Blanket bog	M17a <i>Trichophorum germanicum</i> - <i>Eriophorum vaginatum</i> blanket mire, <i>Drosera rotundifolia</i> - <i>Sphagnum</i> spp. sub-community	222.99/47.54%
E1.7	Wet modified bog	M17a <i>Trichophorum germanicum</i> - <i>Eriophorum vaginatum</i> blanket mire, <i>Drosera rotundifolia</i> - <i>Sphagnum</i> spp. sub-community	34.65%/7.39
E2.1	Acid/neutral flush	M6c <i>Carex echinata</i> - <i>Sphagnum recurvum</i> / <i>auriculatum</i> mire, <i>Juncus effuses</i> sub-community	14.98/3.19%
I2.1	Quarry	NA	0.25/0.05%
			Total Area: 469